

## Math 3342 Exam #3 practice

These are sample questions, but this is not meant to be exhaustive of everything that might be on the final exam.

**Question 1.** Please make a Turing machine that accepts any string of the form  $axa$  for  $x \in \{a, b\}^*$ , and runs forever when the input is not of this form.

**Question 2.** Let  $M$  and  $N$  be these machines using the alphabet  $\Sigma = \{a, b\}$ :



(These each are machines of a single starting state with no transition arrows.)

- Is  $M$  legal when interpreted as a DFA? If so, say what  $L(M)$  is. If not, say why not.
- Is  $N$  legal when interpreted as a DFA? If so, say what  $L(N)$  is. If not, say why not.
- Is  $M$  legal when interpreted as a NFA? If so, say what  $L(M)$  is. If not, say why not.
- Is  $N$  legal when interpreted as a NFA? If so, say what  $L(N)$  is. If not, say why not.
- Is  $M$  legal when interpreted as a Turing Machine? If so, say what  $L(M)$  is. If not, say why not.
- Is  $N$  legal when interpreted as a Turing Machine? If so, say what  $L(N)$  is. If not, say why not.

**Question 3.** For this problem, let  $G$  be this grammar:

$$\begin{aligned} S &\rightarrow aTX \\ T &\rightarrow bR \\ R &\rightarrow S \mid \varepsilon \\ X &\rightarrow aX \mid \varepsilon \end{aligned}$$

- Please give a grammar derivation for some nonempty string in  $L(G)$ .
- Please describe in words or set theory notation the set  $L(G)$ .
- Please give a regular expression equivalent to  $G$ , or explain why this is not possible.
- Please give a stack machine equivalent to  $G$ , and show a stack derivation of some specific nonempty string.

**Question 4.** Please make a Turing machine with language  $\{a^{2^n}cb^n\}$ .

**Question 5.** Please make a Turing machine that computes this function:  $f(a^n) = a^{n \bmod 2}$ . (The answer left on the tape should be either  $a$  or  $\varepsilon$ , depending if  $n$  is even or odd.)

**Question 6.** Please give an NFA for this regular expression:  $(a + b)^*ab + a^*b$

**Question 7.** In each part, either show that the language is not regular, or show that it is regular.

a)  $\{a^n b x a b^n \mid x \in \{a, b\}^*\}$

b)  $\{a^n x b^m \mid x \in \{a, b\}^*\}$


**Question 8.** In each part, please give a grammar for each language:

a)  $\{ab^n a\}$

b)  $\{a^n b^m c^{2m}\}$


c)  $\{a^n x (ab)^n \mid x \in \{a, b\}^*\}$

**Question 9.** Please convert this NFA to an equivalent DFA:



## Answers!

1.



2. a)  $M$  is not a legal DFA, since it would need outgoing arrows for each letter in the alphabet.  
 b)  $N$  is not a legal DFA for the same reason.  
 c)  $M$  is a legal NFA. Its language is  $L(M) = \emptyset$ .  
 d)  $N$  is a legal NFA. Its language is  $L(N) = \{\epsilon\}$ .  
 e)  $M$  is a legal Turing Machine, with  $L(M) = \emptyset$ .  
 f)  $N$  is a legal Turing Machine, with  $L(N) = \{a, b\}^*$ .

3. a) I'll derive  $ab$ :  $S \Rightarrow aTX \Rightarrow abRX \Rightarrow abX \Rightarrow ab$

b)  $L(G) = \{(ab)^n a^m \mid n \geq 1, m \geq 0\}$

c)  $ab(ab)^*a^*$


d) My stack machine:

read	pop	push
$\epsilon$	$S$	$aTX$
$\epsilon$	$T$	$bR$
$\epsilon$	$R$	$S$
$\epsilon$	$R$	$\epsilon$
$\epsilon$	$X$	$aX$
$\epsilon$	$X$	$\epsilon$
$a$	$a$	$\epsilon$
$b$	$b$	$\epsilon$

A derivation of  $ab$ :


$$(ab, S) \mapsto (ab, aTX) \mapsto (b, TX) \mapsto (b, bRX) \mapsto (\epsilon, RX) \mapsto (\epsilon, X) \mapsto (\epsilon, \epsilon)$$

4. My strategy is to mark 2  $a$ s, then go right and mark one  $b$ . Make sure not to loop on  $c/c$ , since this would allow for any number of  $c$ 's. We want to require just one.




5. This one is a bit tricky. My strategy is: we see an  $a$  first, then move R and look for another  $a$ . If there isn't one, then we had only one  $a$  left, so the string is odd, so leave the  $a$  and stop. If we see a second  $a$ , then back up and blank both  $a$ 's. Then repeat.


This ends up blanking 2  $a$ s at a time, so we end up with either empty, or one leftover  $a$ , depending if  $n$  was even or odd.



Here's another solution which is cute but a bit cryptic:



6.



7. a) This one is not regular: let  $D_i = \frac{d}{da^i} L = \{a^{n-i} b x a b^n \mid x \in \{a, b\}^*\}$ . Then these are all different, so  $L$  has infinitely many derivatives, so  $L$  is not regular.

b) This one is regular. Here is a regular expression:  $a^*(a + b)^*b^*$ .

8. a)

$$S \rightarrow aTa$$

$$T \rightarrow bT \mid \epsilon$$

b)

$$S \rightarrow AT$$

$$A \rightarrow aA \mid \epsilon$$

$$T \rightarrow bTcc \mid \epsilon$$

c)

$$S \rightarrow aSab \mid T$$

$$T \rightarrow aT \mid bT \mid \epsilon$$

9.

